

The Optical Gravitational Lensing Experiment. Search for Planetary and Low-Luminosity Object Transits in the Galactic Disk. Results of 2001 Campaign – Supplement*

A. Udalski¹, K. Żebruń¹, M. Szymański¹,
M. Kubiak¹, I. Soszyński¹, O. Szewczyk¹,
L. Wyrzykowski¹, and G. Pietrzyński^{2,1}

¹Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warszawa, Poland
e-mail:

(udalski,zebrun,msz,mk,soszynsk,szewczyk,wyrzykow,pietrzyn)@astrouw.edu.pl

² Universidad de Concepción, Departamento de Física, Casilla 160-C,
Concepción, Chile

ABSTRACT

The photometric data collected during 2001 season OGLE-III planetary/low luminosity object transit campaign were reanalyzed with the new transit search technique – the BLS method by Kovacs, Zucker and Mazeh. In addition to all presented in our original paper transits, additional 13 objects with transiting low-luminosity companions were discovered. We present here a supplement to our original catalog – the photometric data, light curves and finding charts of all 13 new objects.

The model fits to the transit light curves indicate that a few new objects may be Jupiter-sized ($R < 1.6 R_{\text{Jup}}$). OGLE-TR-56 is a particularly interesting case. Its transit has only 13 mmag depth, short duration and a period of 1.21190 days. Model fit indicates that the companion may be Saturn-sized if the passage were central.

Spectroscopic follow-up observations are encouraged for final classification of the transiting objects as planets, brown dwarfs or late M-type dwarf stars.

We also provide the most recent ephemerides of other most promising planetary transits: OGLE-TR-10 and OGLE-TR-40 based on observations collected in June 2002.

All photometric data are available to the astronomical community from the OGLE Internet archive.

1. Introduction

In the paper by Udalski *et al.* (2002), the OGLE photometric survey presented results of the pilot campaign aiming at discovery of small depth transits caused by planets or low-luminosity small objects like brown dwarfs or late type M-dwarfs. The campaign turned out to be very successful – 46 objects with transiting objects causing small (< 0.08 mag) box-shaped drop of brightness were detected. Simple transit model assuming completely dark transiting object and

*Based on observations obtained with the 1.3 m Warsaw telescope at the Las Campanas Observatory of the Carnegie Institution of Washington.

limb darkening of the primary star was fitted to the light curves of transits, indicating small sizes of transiting companions. In several cases the results indicated Jupiter-sized objects ($R < 1.6 R_{\text{Jup}}$).

Unfortunately, the photometry alone cannot unambiguously distinguish between Jupiter size planets and other low-luminosity objects: brown dwarfs and late type M dwarfs. All of them have radii of the order of $0.1\text{--}0.2 R_{\odot}$ ($1\text{--}2 R_{\text{Jup}}$). Thus, a spectroscopic follow-up and a measurement of the radial velocity amplitude of the stars is needed to determine the masses of transiting companions and final classification.

Large number of discovered objects with transiting companions makes the transit method of planetary search very attractive as the efficiency of search can be much higher than that of standard spectroscopic searches. Moreover, non-planetary companions like brown dwarfs or faint late M-type dwarf are also very interesting astronomically and their parameters are poorly known. It should be noted that photometry of transits combined with precise spectroscopy is the only method of unambiguous determination of all basic parameters like dimensions and masses.

Recently, Kovacs, Zucker and Mazeh (2002) proposed a new method of analysis of large photometric datasets for transit detection: the Box-fitting Least Squares (BLS) algorithm. Encouraged by promising simulations on artificial datasets presented by Kovacs *et al.* (2002) we ran the method on photometric data of all objects presented in Udalski *et al.* (2002). All of them were easily found by the BLS method. Encouraged even more, we decided to run the algorithm on photometric data of all $\approx 52\,000$ Galactic disk stars selected for transit search. Results were very impressive – 13 additional objects with transiting low-luminosity companions were detected by the BLS method, increasing the total number of transit objects detected in the 2001 OGLE-III campaign data to 59.

This paper is a supplement to the original paper presenting results of the 2001 OGLE-III planetary/low-luminosity object transit campaign (Udalski *et al.* 2002). We show here photometry, light curves and finding charts as well as results of preliminary analysis of transit light curves for all 13 new objects detected with the BLS method. Photometry of new OGLE-III stars with transits is available to the astronomical community from the OGLE Internet archive.

2. Observational Data

Observations were collected with the 1.3-m Warsaw telescope at the Las Campanas Observatory, Chile, (operated by the Carnegie Institution of Washington) equipped with a large field CCD mosaic camera, consisting of eight 2048×4096 pixel SITe ST002A detectors. The pixel size of each of the detectors is $15\,\mu\text{m}$ giving the $0.26\,\text{arcsec/pixel}$ scale at the focus of the Warsaw telescope. Full field of view of the camera is about $35 \times 35.5\,\text{arcmins}$. The gain of each chip is adjusted to be about $1.3\,\text{e}^-/\text{ADU}$ with the readout noise from about 6 to $9\,\text{e}^-$, depending on the chip.

The photometric data were obtained during 32 nights spanning 45 days from June 12, 2001. Three fields located in the densest stellar regions in the direction of the Galactic center were monitored continuously up to 35 times per night. After that period single observations of these fields were done once every few nights till the end of 2001 Galactic bulge season in October 2001. Almost all observations were made in the *I*-band filter. The exposure time of each image was set to 120 seconds. Altogether about 800 epochs were collected for each field during the 2001 season.

3. Search for Transits with the BLS Method

About 52 000 Galactic disk stars with accuracy of photometry (*rms*) better than 0.015 mag were selected from about 5 million stars observed in three fields during the 2001 OGLE-III campaign. Details of the selection procedure are described in Udalski *et al.* (2002). All these objects were subject to the BLS algorithm for transit search.

We wrote the BLS code procedure based on the description and exemplary Fortran code provided by Kovacs *et al.* (2002). We used the algorithm with 200 phase bins, and the BLS statistics was calculated for frequencies from 0.1 [1/day] ($P=10$ days) to 0.95 [1/day] ($P=1.053$ days) with the step of 0.0001 [1/day]. Transits of duration from 0.015 to 0.15 in phase were searched for.

Only objects for which the highest peak in the BLS power spectrum (BLS statistics *vs.* frequency) exceeded the effective signal-to-noise parameter $\alpha=9.0$ and signal detection efficiency $SDE=3.0$ (see definitions in Kovacs *et al.* 2002) were further analyzed. However, for the lowest signal-to-noise detection we required proportionally larger SDE value to avoid too many false detections. Linear relation α *vs.* SDE serving as a detection threshold was experimentally established based on results of the BLS run on the transit sample from Udalski *et al.* (2002).

It should be noted that the BLS method is very efficient numerically. It took only several CPU hours to run all 52 000 stars through the BLS algorithm compared to weeks with the method used in the original search by Udalski *et al.* (2002).

4. Results of the BLS Search

Beside of all transit objects detected in Udalski *et al.* (2002), the BLS search detected 13 additional objects with low-luminosity transiting objects. Actually, the number of new detections was larger. However, on the final list we only left the cases that light curves indicate transit with high probability. Objects with shallow but clearly triangle-shaped eclipses were removed as they are probably grazing eclipses of normal stars. Also objects with indications of secondary eclipses – very likely, regular eclipsing stars blended with brighter disk stars were not included.

We present new transit objects in identical form as for the original detections: in the form of a catalog and atlas of light curves. Table 1 contains basic data for each star: abbreviation in the form OGLE-TR-NN, equatorial coordinates (J2000.0 epoch), orbital period, epoch of mid-eclipse, I -band magnitude and an approximate $V - I$ color outside eclipse, the depth of eclipse, number of transits observed (N_{tr}), and remarks. OII acronym in the remarks column indicates that the object was observed in the OGLE-II phase of the OGLE survey. For these stars both – the mean I -band magnitude and $V - I$ color from OGLE-II photometric maps (Udalski *et al.* 2002, in preparation) were used to set the zero point of photometry, and both values are accurate to better than 0.05 mag. In the remaining cases the accuracy of the magnitude scale and $V - I$ color is not better than 0.1 mag. It should be also noted that in the case of OGLE-TR-56 its ephemeris was updated based on observations collected in 2002 observing season up to June 2002.

Table 1
OGLE-III planetary and low luminosity object transits

Name	RA (J2000)	DEC (J2000)	P [days]	T_0 -2452000	I [mag]	$V - I$ [mag]	ΔI [mag]	N_{tr}	Rem.
OGLE-TR-47	17 ^h 52 ^m 15 ^s 53	-30°13'54''0	2.33590	76.45988	15.602	1.26	0.019	7	OII
OGLE-TR-48	17 ^h 51 ^m 17 ^s 11	-30°03'01''3	7.22550	74.19693	14.770	1.10	0.022	2	
OGLE-TR-49	17 ^h 51 ^m 14 ^s 41	-29°54'23''8	2.69042	75.53363	16.180	1.72	0.034	2	
OGLE-TR-50	17 ^h 53 ^m 48 ^s 08	-29°56'00''9	2.24861	76.26084	15.923	1.48	0.048	3	OII
OGLE-TR-51	17 ^h 52 ^m 36 ^s 00	-29°37'28''9	1.74832	74.99241	16.714	1.42	0.030	4	OII
OGLE-TR-52	17 ^h 52 ^m 54 ^s 01	-29°46'33''8	1.32565	73.84778	15.593	1.41	0.034	8	OII
OGLE-TR-53	17 ^h 53 ^m 09 ^s 82	-30°06'29''6	2.90565	77.59653	15.998	1.13	0.034	4	OII
OGLE-TR-54	17 ^h 52 ^m 57 ^s 52	-30°05'33''1	8.16256	74.73849	16.473	1.82	0.026	3	OII
OGLE-TR-55	17 ^h 57 ^m 28 ^s 50	-29°43'50''4	3.18473	77.04692	15.800	1.48	0.019	4	
OGLE-TR-56	17 ^h 56 ^m 35 ^s 51	-29°32'21''2	1.21190	72.68492	15.300	1.26	0.013	11	
OGLE-TR-57	17 ^h 55 ^m 53 ^s 20	-29°22'28''7	1.67483	75.46493	15.698	1.51	0.026	5	OII
OGLE-TR-58	17 ^h 55 ^m 08 ^s 27	-29°48'51''3	4.34244	76.81982	14.751	1.20	0.030	2	OII
OGLE-TR-59	17 ^h 55 ^m 03 ^s 30	-29°48'48''4	1.49709	73.15416	15.196	1.22	0.026	9	OII

Light curves in graphical form are presented in Appendix: full phased light curve and close up around the eclipse. Additionally we also provide a finding chart – 60×60 arcsec subframe of the I -band reference image centered on the star. North is up and East to the left in these images.

5. Discussion

The BLS method of transit detection (Kovacs *et al.* 2002) proved to be a very effective and efficient tool for analysis of huge photometric datasets for peri-

odic transits. Application of the method to the real photometric data collected during the 2001 OGLE-III transit campaign allowed to increase significantly the sample of low-luminosity transit objects, which is now totaling at 59. It is worth noting that the BLS method detects transit with smaller depth – the smallest depth in the case of the OGLE-TR-56 transit is only 13 mmag – and also transits in more noisy data (fainter objects).

For all new transit objects we performed an analysis of transit light curve similar as in our original paper (Udalski *et al.* 2002). We estimated sizes of transiting objects by modeling their light curves assuming a completely dark companion and using formulae provided by Sackett (1999). These were numerically integrated at the appropriate phase points to produce model light curves. Details can be found in Udalski *et al.* (2002).

The smallest size of the star and its companion is obtained when the transit is central, *i.e.*, for the inclination $i = 90^\circ$. Non-central passage (smaller i) requires a larger size of both, the star and its companion. Table 2 lists the minimum size of the transiting companion and the corresponding size of the star (assuming $M_s = 1 M_\odot$) to provide information on the expected dimensions of the transiting objects. It should be remembered that semi-major axis of the orbit scales as $M^{1/3}$, and therefore the same scaling is appropriate for the sizes of stars and companions listed in Table 2. In the close-up windows in the Appendix we show model light curves for central transit ($i = 90^\circ$) drawn with a continuous bold line.

T a b l e 2
Dimensions of stars and companions for central passage ($M_s = 1 M_\odot$)

Name	R_s [$R_\odot$]	R_c [$R_\odot$]
OGLE-TR-47	1.95	0.23
OGLE-TR-48	1.21	0.16
OGLE-TR-49	0.97	0.15
OGLE-TR-50	1.16	0.22
OGLE-TR-51	0.85	0.13
OGLE-TR-52	1.27	0.20
OGLE-TR-53	0.76	0.12
OGLE-TR-54	1.38	0.19
OGLE-TR-55	0.93	0.11
OGLE-TR-56	0.80	0.08
OGLE-TR-57	1.85	0.26
OGLE-TR-58	1.75	0.26
OGLE-TR-59	1.15	0.16

Table 2 indicates that in several cases the new transiting companions are of Jupiter size ($< 1.6 R_{\text{Jup}}$). They may be planets, or brown dwarfs or low-luminosity M-type stars. The spectroscopic follow-up will provide necessary

clarification.

The most intriguing object within this group is OGLE-TR-56. Table 2 indicates that the size of the transiting companion is smaller than the Saturn size if its transit is central. However, even for inclination as low as $i = 83^\circ$, its radius is still equal to one Jupiter radius. This object is certainly very small. The color of the primary star indicates a star of smaller size than the Sun consistent with Table 2. If confirmed spectroscopically to be a planet, OGLE-TR-56 would have the shortest orbital period (1.21190 days) and would be one of the smallest extrasolar planets.

Alternatively the transiting object in OGLE-TR-56 might be a brown dwarf. Recently, Santos *et al.* (2002) reported possible detection of a brown dwarf around HD 41004 with a period of only 1.3 days. Also, possible detection of a brown dwarf in the short period Hyades binary system RHy 403 was reported by Reid and Mahoney (2000). OGLE-TR-56 could be another example indicating that the so called “brown dwarf desert” – lack of brown dwarfs with short periods on small orbits, is not that empty as generally thought.

The least interesting alternative would be a blend of a regular Galactic bulge eclipsing star with a bright disk star which could mimic the low depth transit. However, this case is not very likely – the shape of the transit light curve, namely short time of descending and ascending branches of the transit compared to long phase of totality, makes this explanation less probable.

It is worth adding here that two other most promising planetary transit objects: OGLE-TR-40 and OGLE-TR-10, discovered in the original paper (Udalski *et al.* 2002), were observed for expected transits on a few nights in June 2002. The main goal of these observations was the improvement of ephemerides that due to short time baseline in 2001 could be inaccurate after longer period of time. For both objects a few transits were covered and the new best ephemerides, based now on more than one hundred orbital cycles, are:

$$\text{HJD}_{\min} = 2452060.03667 + 3.43079 \cdot E \quad (\text{OGLE} - \text{TR} - 40)$$

$$\text{HJD}_{\min} = 2452070.21900 + 3.10140 \cdot E \quad (\text{OGLE} - \text{TR} - 10)$$

The photometric data of new OGLE-III transit objects presented in this paper are available in the electronic form from the OGLE archive:

<http://www.astroww.edu.pl/~ogle>

ftp://ftp.astroww.edu.pl/ogle/ogle3/transits/new_2001_transits

or its US mirror

<http://bulge.princeton.edu/~ogle>

ftp://bulge.princeton.edu/ogle/ogle3/transits/new_2001_transits

Acknowledgements. We would like to thank Prof. B. Paczyński for many interesting discussions and comments. The paper was partly supported by the Polish KBN grant 2P03D01418 to M. Kubiak. Partial support to the OGLE project was provided with the NSF grant AST-9820314 and NASA grant NAG5-12212 to B. Paczyński.

REFERENCES

- Kovacs, G., Zucker, S., and Mazeh, T. 2002, *Astron. Astrophys.*, in press; astro-ph/0206099.
- Sackett, P.D. 1999, in: "Planets Outside the Solar System: Theory and Observations," Eds. J.-M. Mariotti and D. Alloin, NATO-ASI Series, Kluwer, 189.
- Reid, I.N., and Mahoney, S. 2000, *MNRAS*, **316**, 827.
- Santos, N.C. *et al.* 2002, *Astron. Astrophys.*, in press; astro-ph/0206213.
- Udalski, A., Paczyński, B., Żebruń, K., Szymański, M., Kubiak, M., Soszyński, I., Szewczyk, O., Wyrzykowski, L., and Pietrzyński, G. 2002, *Acta Astron.*, **52**, 1.

This figure "app_1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0207133v1>

This figure "app_2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0207133v1>

This figure "app_3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0207133v1>

This figure "app_4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0207133v1>

This figure "app_5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0207133v1>

This figure "app_6.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0207133v1>

This figure "app_7.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0207133v1>